

Weather conditions for wildfires have changed in Southern Europe

August 19th, 2021



Wildre in Portugal in 2014 (photo: Graeme Darbyshire, www. jckr.com)

The Mediterranean is Europe's wildftre hotspot region. Mediterranean wildftres generate 85% of the total burnt area of Europe, annually.

Extreme ftre weather conditions in the European part of the Mediterranean have undergone significant changes in the past 30-years. These changes have been studied for the period 1987-2016. According to the authors of this study, this is the first high-resolution analysis of near-present ftre weather covering the entire European part of the Mediterranean.

Fire Weather Index

Measurement data is often too scarce or inconsistent over a sufficiently long period to be used for properly assessing fire weather. Computer models can be used, however, to provide the spatially and temporally complete datasets required for characterizing fire weather and detecting trends. For this study, a high-resolution dataset of fire weather conditions was computed from model simulations. These fire weather conditions are air temperature, relative humidity, wind speed, and precipitation. These variables can be combined into an

index called the Fire Weather Index, an indicator of potential fire intensity. The index includes a rating of the sammability (moisture content) of surface litter and organic material in deeper soil layers.

The Fire Weather Index was calculated for the entire study area throughout the 30-year period, and this data was analysed for spatial patterns and temporal trends.

Spatial patterns

Fire weather conditions are generally most extreme in both the western and eastern parts of the Euro-Mediterranean. In addition, the fire weather index increases from north to south. Adverse fire weather conditions occur most frequently in Southern Spain, South-eastern Continental Greece, South Aegean Sea islands (Cyclades, Dodecanese, Crete), Eastern and Central Turkey, and Cyprus.

Fire season

For the entire Euro-Mediterranean, critical ftre weather conditions typically occur from May through September. In the Southern Euro-Mediterranean, the Fire Weather Index peaks around mid to end of July, when temperature reaches its annual maximum, and relative humidity and precipitation reach their annual minimum. In the Northern Euro-Mediterranean, the index peaks later in summer (mid of August to start of September).

Trends

In general, there is a trend towards a warmer and drier climate in this part of Europe that promotes extreme fire weather conditions. There are spatial differences, however. The analysis indicates increasing trends in the occurrence of extreme fire weather over Portugal and Spain (the Iberian Peninsula) and eastern Balkans. Decreasing trends have been found over the Southeast Mediterranean, particularly over Southern Italy and Greece. Not surprisingly, these trends are driven by changes towards drier or wetter conditions, respectively.

Over the Iberian Peninsula and eastern Balkans (around the Black Sea), the number of days with adverse fire weather conditions has increased by 0.75 to 1 day per year, on average, during the 1987-2016 period. On the other hand, this number of days has decreased by 0.75 to 1 day per year during the same period in parts of Southern Greece, Cyprus, and Southern Turkey.

Source: Giannaros et al., 2020. International Journal of Climatology.